

SPIN, ACCRETION AND THE COSMOLOGICAL GROWTH OF SUPERMASSIVE BLACK HOLES

STUART L. SHAPIRO^{1,2}

¹ Department of Physics, University of Illinois at Urbana-Champaign,
 Urbana, IL 61801-3080 and

² Department of Astronomy and NCSA, University of Illinois at Urbana-Champaign,
 Urbana, IL 61801-3080

Draft version February 2, 2008

ABSTRACT

If supermassive black holes (SMBHs) are the energy sources that power quasars and active galactic nuclei, then QSO SDSS 1148+5251, the quasar with the highest redshift ($z_{\text{QSO}} = 6.43$), hosts a supermassive black hole formed within ~ 0.9 Gyr after the Big Bang. This requirement places constraints on the cosmological formation of SMBHs, believed to grow from smaller initial seeds by a combination of accretion and mergers. We focus on gas accretion onto seeds produced by the collapse of Pop III stars at high redshift. We incorporate the results of recent relativistic, MHD simulations of disk accretion onto Kerr black holes to track the coupled evolution of the masses and spins of the holes. We allow for an additional amplification of $\sim 10^4$ in the mass of a typical seed due to mergers, consistent with recent Monte Carlo simulations of hierarchical mergers of cold, dark matter halos containing black hole seeds. We find that the growth of Pop III black hole remnants to $\sim 10^9 M_\odot$ by $z_{\text{QSO}} \gtrsim 6.43$ favors MHD accretion disks over standard thin disks. MHD disks tend to drive the holes to a submaximal equilibrium spin rate $a/M \sim 0.95$ and radiation efficiency $\epsilon_M \sim 0.2$, while standard thin disks drive them to maximal spin ($a/M = 1$) and efficiency ($\epsilon_M = 0.42$). This small difference in efficiency results in a huge difference in mass amplification by accretion at the Eddington limit. The MHD equilibrium efficiency is consistent with the observed ratio of the QSO plus AGN luminosity density to the local SMBH mass density. Our prototype analysis is designed to stimulate the incorporation of results from relativistic stellar collapse and MHD accretion simulations in future Monte Carlo simulations of hierarchical structure formation to better determine the cosmological role of SMBHs and their mass and spin distributions.

Subject headings: black hole physics—accretion—MHD—cosmology—quasars

1. INTRODUCTION

There is substantial evidence that supermassive black holes (SMBHs) with masses in the range $10^6 - 10^{10} M_\odot$ exist and are the engines that power active galactic nuclei (AGNs) and quasars (Rees 1984, 1998, 2001; Macchetto 1999). There is also ample evidence that SMBHs reside at the centers of many, and perhaps most, galaxies (Richstone et al. 1998; Ho 1999), including the Milky Way (Genzel et al. 1997; Ghez et al. 2000, 2003; Schödel et al. 2002).

The highest redshift of a quasar discovered to date is $z_{\text{QSO}} = 6.43$, corresponding to QSO SDSS 1148+5251 (Fan et al. 2003). Accordingly, if they are the energy sources in quasars (QSOs), the first supermassive black holes must have formed prior to $z_{\text{QSO}} = 6.43$, or within $t = 0.87$ Gyr after the Big Bang in the concordance Λ CDM cosmological model. This requirement sets a significant constraint on black hole seed formation and growth mechanisms in the early universe. Once formed, black holes grow by a combination of mergers and gas accretion. An important clue to the growth process is provided by the ratio R of the observed QSO plus AGN luminosity density to the local SMBH mass density, since R is related to the mean radiative efficiency ϵ_M of accretion onto black holes (Soltan 1982): $\epsilon_M \geq R$. Recent measurements suggest $0.1 \lesssim R \lesssim 0.2$ (Yu & Tremaine 2002; Elvis, Risaliti & Zamorani 2002), a range consistent with disk accretion onto black holes. Another clue is provided the estimated ratios ϵ_L of the bolometric-

to-Eddington luminosities of the broad-line quasars in a Sloan Digital Sky Survey sample of 12,698 quasars in the redshift interval $0.1 \leq z \leq 2.1$. This survey supports the value $\epsilon_L \approx 1$ as a physical upper limit (McLure & Dunlop 2004). Barring an extraordinary coincidence, the range of values inferred from observations for both ϵ_M and ϵ_L suggest that a significant fraction of the mass of SMBHs is acquired by gas accretion. Together, these two parameters control the rate of growth of black holes by accretion and are crucial in determining whether or not initial seed black holes formed at high redshift have sufficient time to grow to SMBHs to explain quasars and AGNs.

The more massive the initial seed, the less time is required for it to grow to SMBH scale and the easier it is to have a SMBH in place by $z \geq 6.43$. One possible progenitor that readily produces a SMBH is a supermassive star (SMS) with $M \gg 10^3 M_\odot$ (see, e.g., Shapiro 2004 for a recent review and references). SMSs can form when gaseous structures build up sufficient radiation pressure to inhibit fragmentation and prevent normal star formation; plausible cosmological scenarios have been proposed that can lead to this occurrence (Gnedin 2001; Bromm & Loeb 2003). SMSs supported by radiation pressure will evolve in a quasi-stationary manner to the point of onset of dynamical collapse due to a general relativistic radial instability (Chandrasekhar 1964a,b; Feynman, unpublished, as quoted in Fowler 1964). The collapse of a nonrotating, marginally unstable, spherical SMS of

mass M leads directly to the formation of a nonrotating Schwarzschild black hole of the same total mass (Shapiro & Teukolsky 1979). But like most stars formed in nature, SMSs will be rotating. In fact, in the event that viscosity and/or turbulent magnetic fields are present to drive these stars to uniform rotation, they are likely to be maximally rotating (i.e. at the mass-shedding limit) by the time they reach the onset of collapse (Baumgarte & Shapiro 1999). Recent relativistic hydrodynamic simulations have shown that unstable, maximally rotating SMSs of arbitrary mass M inevitably collapse to SMBHs of mass $\sim 0.9M$ and spin parameter $a/M \sim 0.75$; the rest of the mass goes into an ambient disk about the hole (Shibata & Shapiro 2002; Shapiro & Shibata 2002; Shapiro 2004).

But the fact remains that SMSs have yet to be observed and there is no concrete evidence that they actually form in the early universe. Moreover, simulations of cosmological structure formation performed to date indicate that the first generation of stars are more likely to be zero-metallicity Pop III stars in the range $10^2 - 10^3 M_\odot$ (Bromm, Coppi & Larson 1999, 2002; Abel, Bryan, & Norman 2000, 2002; but note that Norman 2004 reports preliminary indications that supermassive stellar objects with $M \gtrsim 10^4 M_\odot$ may be forming as *second* generation stars at $10 \lesssim z \lesssim 15$ in his latest simulations). So the most conservative hypothesis is that the seed black holes that later grow to become SMBHs originate from the collapse of Pop III stars (Madau & Rees 2001) and not SMSs, and this is the ansatz we shall explore here. Newtonian simulations suggest that Pop III stars with masses in the range $M \sim 60 - 140 M_\odot$ and $M \gtrsim 260 M_\odot$ collapse directly to black holes, while stars with $M \sim 140 - 260 M_\odot$ undergo explosive annihilation via pair-creation processes (Heger et al. 2003). The upper limit to the mass of a Pop III star is set by accretion over a stellar lifetime, yielding $M \lesssim 600 M_\odot$ (Onukai & Palla 2003; Yoshida et al. 2003). Abel et al. (2002) argue that first generation stars significantly larger than $100 M_\odot$ are likely to explode before they have time to accrete to larger masses. The Pistol Star is an example of an existing star believed to have a mass $\gtrsim 200 M_\odot$, although it has high metallicity (Figer et al. 1998). Very massive stars are dominated by thermal radiation pressure, so the catastrophic collapse of those that do not explode will be hydrodynamically similar to the collapse of SMSs, producing black holes with masses comparable to those of their progenitors. The higher the mass of the black hole seed, and the earlier it forms in the universe, the easier it is for it to grow to a SMBH, hence the more conservative will be any constraints imposed on the cosmological black hole growth rate by the existence of a SMBH by $z_{\text{QSO}} = 6.43$. Accordingly, we take the highest range of plausible values for the masses of black hole seeds to establish the most conservative (robust) constraints, adopting $100 \leq M/M_\odot \leq 600$ for the range of black hole seeds arising from the collapse of Pop III stars at $z \leq 40$.

The most detailed studies of SMBH formation to date involve detailed, Monte Carlo simulations that follow the cosmological growth of a distribution of black hole seeds by a combination of discrete, stochastic mergers as well as gas accretion (see, e.g., Haehnelt & Kauffmann 2000; Volonteri, Haardt & Madau 2003; Brom-

ley et al. 2004; Haehnelt 2004; Haiman 2004; Yoo & Miraldi-Escudé 2004; and references therein). Both processes are assumed to take place in the context of the cold dark matter (CDM) model, where dark matter halos merge hierarchically, and black holes are assumed to settle, merge and accrete in their gaseous centers. Typically, the stellar dynamical processes that lead to mergers, as well as the hydrodynamical processes that fuel accretion, are modeled in these analyses by implementing simple, physically plausible, rules rather than by detailed integrations of the governing dynamical and hydrodynamical equations of motion. Performing such “first principles” integrations would prove prohibitive in this context.

In this paper we focus on SMBH cosmological growth by accretion. We identify and explore the main variables that govern this process and ultimately influence the outcome of detailed Monte Carlo simulations that track SMBH growth. We incorporate some of the most recent findings of relativistic magnetohydrodynamical (MHD) simulations of gas accretion onto black holes and explore their cosmological implications for SMBH evolution. We show how the evolution and amplification of black hole *mass* by accretion is intimately tied to the evolution of black hole *spin*, probing some of the implications of our earlier survey and analysis of black hole spin evolution (Gammie, Shapiro & McKinney 2004) for cosmology. We demonstrate how in principle the very existence of a quasar at redshift $z_{\text{QSO}} = 6.43$ can help constrain the formation epoch and/or size of black hole seeds and select among competing models of accretion.

Specifically, recent relativistic MHD simulations predict the radiation efficiency ϵ_M as a function of the BH spin parameter a/M ; they also predict the spin-up rate as a function of a/M . Monte Carlo simulations that determine cosmological SMBH growth must integrate coupled mass and spin evolution prescriptions versus time for each accreting BH to produce reliable growth histories and final SMBH masses and spins. This paper provides prototype integrations of the coupled evolution equations, focussing on the history of the progenitor of SDSS 1148+5251 and demonstrating the significant differences in the outcomes for standard thin disk accretion models versus recent MHD models.

Our discussion is simplified and illustrative at best: much of the input physics involving black hole seed formation, accretion flows, and mergers is still being developed. Our main goal is to isolate some of the underlying local physical issues and parameters pertaining to accretion to better understand their role in determining the global outcome of the cosmological Monte Carlo simulations of SMBH build-up during hierarchical structure formation. Excellent overviews of the input physics have appeared elsewhere (see, e.g. Haiman & Quataert 2004, and references to earlier work). Also, earlier treatments have considered some of the constraints on cosmological growth imposed by the recent discoveries of luminous quasars at high redshift (e.g., Haiman & Loeb 2001; Haiman 2004; Yoo & Miraldi-Escudé 2004). But here we specifically want to illustrate in the simplest fashion how the most recent findings related to relativistic, MHD accretion flows onto spinning black holes have important implications for evolutionary models of the growth of SMBHs in the early universe. We emphasize by concrete example the point made in Gammie et al. (2003)

that tracking the spin as well as the mass of a black holes is necessary to determine its growth (see also Volonteri et al. 2004). We also show that the value of accretion radiation efficiency, ~ 0.1 , adopted in many Monte Carlo simulations may not be entirely consistent with the latest MHD accretion disk modeling. Determining this parameter, on which the outcome of cosmological simulations of SMBH growth depends very sensitively (exponentially!), is coupled to the spin evolution of the hole; both may now be within our grasp via detailed relativistic MHD simulations of BH accretion.

The calculations performed here are prototypical only; our main aim is to motivate the incorporation of these parameters in more detailed Monte Carlo studies and thereby sharpen some of the rules that enter these simulations. We also hope to provide those members of the computational relativistic MHD community who may not be SMBH Monte Carlo cognoscenti a simple means of extracting the essence of the Monte Carlo simulations, particularly the evolutionary consequences of different accretion models.

In Section 2 we set out the basic equations that describe the coupled evolution of black hole mass and spin by accretion. We also summarize in this section the relations we require from the concordance Λ CDM cosmological model. In Section 3 we integrate the coupled evolution equations to track the evolution of black hole mass and spin as functions of time. In Section 4 we apply the results to the cosmological problem and identify some constraints imposed by the existence of a SMBH at $z \geq 6.43$. In Section 5 we summarize briefly and discuss some caveats and areas for further study.

2. BASIC EQUATIONS

Here we assemble the fundamental black hole accretion evolution equations and review the underlying assumptions on which they are based. We then specify a background cosmological model in the which the growth of the black hole to supermassive size occurs.

2.1. Black Hole Mass and Spin Evolution

Define ϵ_M , the efficiency of conversion of rest-mass energy to luminous energy by accretion onto a black hole of mass M , according to

$$\epsilon_M \equiv L/\dot{M}_0 c^2, \quad (1)$$

where $\dot{M}_0$ is the rate of rest-mass accretion and L is the luminosity. Define ϵ_L , the efficiency of accretion luminosity, according to

$$\epsilon_L \equiv L/L_E, \quad (2)$$

where L_E is the Eddington luminosity, given by

$$L_E = \frac{4\pi M \mu_e m_p c}{\sigma_T} \approx 1.3 \times 10^{46} \mu_e M_8 \text{ erg s}^{-1}. \quad (3)$$

Here we assume that the accretion is dominated by normal, baryonic matter and ignore any contribution of collisionless or self-interacting dark matter (but see, e.g., Ostriker 2000 and Balberg & Shapiro 2002 for alternative scenarios). In particular, we assume that the accreting gas consists of fully ionized atoms and that the principal opacity source is Thomson scattering. The quantity μ_e is the mean molecular weight per electron and m_p is

the proton mass. The black hole growth rate must account for the loss of accretion mass-energy in the form of outgoing radiation according to

$$\frac{dM}{dt} = (1 - \epsilon_M) \dot{M}_0. \quad (4)$$

Combining eqns. (1)–(4) we obtain the black hole growth rate,

$$\frac{dM}{dt} = \frac{\epsilon_L (1 - \epsilon_M)}{\epsilon_M} \frac{M}{\tau}, \quad (5)$$

where τ is the characteristic accretion timescale,

$$\tau \equiv \frac{Mc^2}{L_E} \approx 0.45 \mu_e^{-1} \text{ Gyr}, \quad (6)$$

and is independent of M .

The mass accretion efficiency ϵ_M is typically a function of the black hole spin parameter $a/M = J/M^2$. It changes with time as the spin evolves. It is convenient to express the spin evolution in terms of the nondimensional quantity $s = s(a/M)$, defined by

$$s \equiv \frac{d(a/M)}{dt} \frac{M}{\dot{M}_0}. \quad (7)$$

Inserting eqns. (4) and (5) into eqn. (7) yields the evolution equation for the black hole spin,

$$\frac{d(a/M)}{dt} = \frac{\epsilon_L}{\epsilon_M} \frac{s}{\tau}. \quad (8)$$

In general, eqns. (5) and (8) must be integrated simultaneously to determine the mass and spin evolution of the black hole.

Determining $\epsilon_M(a/M)$ and $s(a/M)$, which are needed to integrate eqns. (5) and (8), requires a gas dynamical model for black hole accretion. We shall assume that the gas has sufficient angular momentum to form a disk about the hole and consider two different accretion disk models: (1) a standard, relativistic, Keplerian “thin disk” with “no-torque boundary conditions” at the innermost stable circular orbit (ISCO) (Pringle & Rees 1972; Novikov & Thorne 1973; see Shapiro & Teukolsky 1983 for review and references) and (2) a relativistic, MHD accretion disk that accounts for the presence of a frozen-in magnetic field in a perfectly conducting plasma (De Villiers & Hawley 2003; De Villiers, Hawley & Krolick 2003; Gammie, Shapiro & McKinney 2003; McKinney & Gammie 2004; and references therein). In the MHD model the magnetorotational instability (MRI; Balbus & Hawley 1991) drives magnetic turbulence and provides the necessary torque to remove angular momentum from the gas and drive the inflow. The MHD model is arguably the most realistic model for disk accretion of magnetized plasma onto a black hole. The standard thin disk model provides a simple, analytic, limiting case that is useful as a point of comparison.

In a standard thin disk corotating with the black hole, the energy and angular momentum per unit rest mass accreted by a black hole are the energy and angular momentum of a unit mass at the ISCO, immediately prior to its rapid plunge and capture by the hole:

$$\begin{aligned} \tilde{E}_{\text{ISCO}} &= \frac{r_{\text{ms}}^2 - 2Mr_{\text{ms}} + a\sqrt{Mr_{\text{ms}}}}{r_{\text{ms}}(r_{\text{ms}}^2 - 3M_h r_{\text{ms}} + 2a\sqrt{Mr_{\text{ms}}})^{1/2}} \\ \tilde{l}_{\text{ISCO}} &= \frac{\sqrt{Mr_{\text{ms}}}(r_{\text{ms}}^2 - 2a\sqrt{Mr_{\text{ms}}} + a^2)}{r_{\text{ms}}(r_{\text{ms}}^2 - 3M_h r_{\text{ms}} + 2a\sqrt{Mr_{\text{ms}}})^{1/2}}, \end{aligned} \quad (9)$$

where the ISCO radius r_{ms} is given by

$$r_{\text{ms}} = M\{3 + Z_2 - [(3 - Z_1)(3 + Z_1 + 2Z_2)]^{1/2}\}, \quad (10)$$

where

$$Z_1 \equiv 1 + \left(1 - \frac{a^2}{M^2}\right)^{1/3} \left[\left(1 + \frac{a}{M}\right)^{1/3} + \left(1 - \frac{a}{M}\right)^{1/3}\right], \quad (11)$$

and

$$Z_2 \equiv \left(3\frac{a^2}{M^2} + Z_1^2\right)^{1/2} \quad (12)$$

(see, e.g., Shapiro & Teukolsky 1983, eqns 12.7.17-12.7.18 and 12.7.24). The mass accretion efficiency and spin evolution parameters corresponding to the thin disk model are then given by

$$\epsilon_M = 1 - \tilde{E}_{\text{ISCO}}, \quad (13)$$

$$s = \tilde{l}_{\text{ISCO}} - 2\frac{a}{M}\tilde{E}_{\text{ISCO}} \quad (\text{standard thin disk}). \quad (14)$$

The MHD disk accretion model of Gammie, Shapiro & McKinney (2004) and McKinney & Gammie (2004) is based on a fully relativistic, axisymmetric simulation of a nonradiative, magnetized plasma onto a Kerr-Schild black hole within the MHD approximation. The initial gas configuration is a torus with an inner radius at $r/M = 6$; in the absence of a magnetic field the torus is constructed to be in equilibrium about the hole (Fishbone & Moncrief 1976). The torus is threaded with a poloidal magnetic field initially and evolves with an adiabatic equation of state (EOS) with an adiabatic index $\Gamma = 4/3$ (to model a radiation-dominated, inner-disk EOS). The source of viscosity is MHD turbulence driven by the MRI instability. The simulations are performed for various black hole spin parameters a/M , holding the value of the spin parameter fixed during the simulation. The evolution proceeds for many dynamical timescales M , until a crude steady-state, with fluctuations, is achieved.

The results of the numerical simulations suggest that in steady-state the radiation efficiency parameter ϵ_M as a function of a/M is remarkably close to the function characterizing the standard thin disk (eqn. 13), even though there is no sharp transition in the surface density at or near the ISCO. However, the spin evolution parameter $s(a/M)$ is different and can be represented reasonably well by the least squares linear fit

$$s = 3.14 - 3.30\frac{a}{M} \quad (\text{MHD disk}), \quad (15)$$

(see Table 2 in McKinney & Gammie 2004). The numerical simulations demonstrate that the above parameters describing steady-state, MHD accretion-disk behavior are not particularly sensitive to the initial conditions in the disk (e.g. the initial B -field). As McKinney & Gammie (2004) discuss, the key results are also quite comparable to those found by De Villiers et al. (2004), who used a different numerical method and took the adiabatic index of the gas to be $\Gamma = 5/3$ instead of $\Gamma = 4/3$. We therefore model a relativistic MHD accretion disk by adopting eqns (13) and (15) in our evolution equations.

The s vs. a/M curve for the MHD disk is roughly parallel to, but somewhat below, the curve for the standard disk (see Fig 4 in Gammie, Shapiro & McKinney 2004 for

a comparison). In particular, the parameter s never falls to zero for a standard thin disk until the hole is maximally rotating at $a/M = 1$, while for an MHD disk s crosses zero at $a/M \approx 0.95$. The crucial physical consequence is that steady accretion via a standard thin disk always causes the central black hole to spin up until it is maximally rotating (Bardeen 1970), while accretion via an MHD disk drives the black hole to spin equilibrium at $a/M \approx 0.95$. While the value of a/M at equilibrium is not determined precisely and may depend on details of the flow geometry (and the presence of radiative cooling), its departure from maximal is robust and is typically $\gtrsim 0.05$ below unity. The difference between $a/M = 1$ and $a/M = 0.95$ is quite substantial physically: for example, the efficiency parameter is $\epsilon_M = 1 - 1/3^{1/2} = 0.42$ for $a/M = 1$, while it is considerably smaller, $\epsilon_M = 0.19$, for $a/M = 0.95$. As we will see below, this difference will have significant consequences for the cosmological growth of a black hole from an initial seed. For comparison we note that the radiative efficiency for disk accretion onto a nonrotating Schwarzschild black hole ($a/M = 0$) is $\epsilon_M = 1 - (8/9)^{1/2} = 0.057$.

If we assume that the supply of gas remains sufficiently copious in the vicinity of the black hole, then the accretion luminosity is likely to be Eddington-limited and nearly constant, with $\epsilon_L \approx 1$. Accretion models characterized by super-Eddington luminosities with $\epsilon_L > 1$ are possible theoretically (Ruszkowski & Begeleman 2003). However, the estimated Eddington ratios of the bolometric luminosities of the broad-line quasars in a Sloan Digital Sky Survey sample of 12,698 quasars in the redshift interval $0.1 \leq z \leq 2.1$ support the value $\epsilon_L \approx 1$ as a physical upper limit (McLure & Dunlop 2004). (Here the mass of the central black hole is estimated from a virial relation, assuming that the gas motions in the broad-line region of the quasars, measured by $H\beta$ and MgII emission lines, are virialized.) The maximum allowed value of ϵ_L establishes the maximum growth rate of a black hole from accretion (see eqn. 5) and, as a result, the most robust (least stringent) constraints on any gas accretion scenario for the growth of a supermassive black hole from a smaller initial seed. To establish these constraints we therefore shall set $\epsilon_L = 1$ in many of our numerical estimates below.

As a point of reference, and for later application, it is useful to integrate eqn. (5) assuming that the mass and luminosity efficiencies both remain constant with time, yielding

$$M(t)/M(t_i) = \exp \left[\frac{\epsilon_L(1 - \epsilon_M)}{\epsilon_M} \frac{(t - t_i)}{\tau} \right], \quad (\epsilon_M, \epsilon_L \text{ constant}), \quad (16)$$

where t_i is the initial time at which the black hole has a mass M_i . Note that the right hand side of eqn. (16) is independent of black hole mass. This fact makes it possible, under certain plausible conditions that we shall specify, to disentangle and track separately the amplification of black hole mass by accretion from the amplification by discrete mergers. Let $M_n(t)$ be the mass of the black hole at time t following its n^{th} merger with another hole at time t_n , where $t_n \leq t \leq t_{n+1}$. Assume that the duration of a merger, as well as the time required for accretion to drive the merged remnant to spin equilibrium (see eqn. 23 below), are both much shorter

than the time interval between mergers, and that the hole continues to accrete steadily throughout this interval. Note that black hole mergers can completely eject black holes from halo centers owing to gravitational wave recoil, and thereby turn off accretion altogether (see discussions of black hole recoil in halos in, e.g., Hut & Rees 1992; Merritt et al. 2004; Madau & Quataert 2004). However, incorporating the most recent recoil calculations (Favata, Hughes & Holz 2004) into simple models of dark halo mergers, Yoo & Miralda-Escudé (2004) conclude that the kick velocities are not sufficiently large to impede black hole growth significantly (cf. Haiman 2004). Note also that a major merger between two black holes of comparable mass may change both the magnitude and direction of the spin of the resulting black hole remnant. Following such a merger, the orientation of the black hole spin may not be aligned with the orientation of the asymptotic gaseous disk at radii $r \gg 100M$ outside the hole. (Rees 1978 and Natarajan & Pringle 1998 point out that the black hole exerts a torque on the asymptotic disk, which also exerts a torque back on the hole, eventually forcing their mutual alignment, but on a timescale that is still uncertain). Moreover, the orientation of the asymptotic disk will likely fluctuate in time, due to the redistribution of gas following mergers of dark halo cores, galaxy mergers, and the tidal disruptions of passing stars by the central hole. However, near the black hole, at radii from $r \sim M$ to $r \sim 20M$, where the bulk of the disk's gravitational energy is released and the hole-disk interactions are strong, the hole's gravito-magnetic field will exert a force on the disk that, when combined with viscous forces and magnetic fields, will drive the disk down into the hole's equatorial plane, (the "Bardeen-Petterson effect"; Bardeen & Petterson 1975). This effect will maintain the alignment between the axis of the inner disk and the spin axis of the hole. Accretion will subsequently drive the hole to spin equilibrium and restore ϵ_M to its equilibrium value.

For the situation described above we can take ϵ_M to be constant and equal to its value at spin equilibrium throughout most of the lifetime of the hole. Assume further that ϵ_L is also constant, which should be the case if the available gas is sufficiently copious that the accretion is always Eddington-limited, whereby $\epsilon_L \approx 1$. Let f_n be the mass amplification of the hole following its n^{th} merger with another hole: $f_n = M_n(t_n)/M_{n-1}(t_n) > 1$. Then we may use eqn. (16) to calculate the total mass amplification from t_i to t_f according to

$$\begin{aligned} \frac{M_f}{M_i} &= \frac{M_N(t_f)}{M_0(t_i)} = \frac{M_0(t_1)}{M_0(t_i)} \frac{M_1(t_1)}{M_0(t_1)} \frac{M_1(t_2)}{M_1(t_1)} \dots \\ &\dots \frac{M_{N-1}(t_N)}{M_{N-1}(t_{N-1})} \frac{M_N(t_N)}{M_{N-1}(t_N)} \frac{M_N(t_f)}{M_N(t_N)} \\ &= \exp[C(t_1 - t_i)] f_1 \exp[C(t_2 - t_1)] \dots \\ &\dots \exp[C(t_N - t_{N-1})] f_N \exp[C(t_f - t_N)] \\ &= f_1 f_2 \dots f_N \exp[C(t_f - t_i)], \end{aligned} \quad (17)$$

where $C = \epsilon_L(1 - \epsilon_M)/(\epsilon_M \tau)$ and is essentially constant. Comparing eqns. (16) and (17) reveals that the net mass amplification due to accretion can be treated as a single multiplicative factor that is independent of the net amplification factor due to mergers, $f = f_1 f_2 \dots f_N$. Thus, for the simple scenario envisioned here, eqn. (16) can

be applied to determine the growth of a black hole by accretion, even when steady growth by accretion is interrupted and augmented by discrete, stochastic black hole mergers.

2.2. Cosmological Model

To relate the time parameter t appearing in the evolution equations to an observable time-like quantity like the redshift z , we adopt the concordance Λ CDM, spatially flat ($k = 0$), cosmological model. All the free parameters in this model that we shall need for our computations have been measured by now, so that the model is uniquely specified.

The basic evolution equation for the Friedmann-Robertson-Walker expansion parameter $a(t)$ is given by

$$H^2 \equiv \left(\frac{\dot{a}}{a}\right)^2 = H_0^2 \left[\Omega_m^0 \left(\frac{a_0}{a}\right)^3 + \Omega_\Lambda^0 \right], \quad (19)$$

where $H(t)$ is Hubble's constant and the normalized mass density parameter Ω_m^0 and cosmological constant parameter Ω_Λ^0 satisfy the relation

$$\Omega_m^0 + \Omega_\Lambda^0 = 1. \quad (20)$$

In the above equations the sub(super)script "0" denotes the value of a quantity at the current epoch, $z = 0$. Recalling that $a/a_0 = 1 + z$ and substituting eqn. (20) into eqn. (19), we can integrate eqn. (19) to obtain $t = t(z)$:

$$t(z) = \frac{2}{3H_0(1 - \Omega_m^0)^{1/2}} \sinh^{-1} \left[\left(\frac{1 - \Omega_m^0}{\Omega_m^0} \right) \frac{1}{(1 + z)^3} \right]^{1/2}. \quad (21)$$

To evaluate $t(z)$ appearing above we adopt the WMAP values for the cosmological parameters: $\Omega_m^0 \approx 0.27$ and $H_0 \equiv 100h$ km/s/Mpc with $h \approx 0.71$ (Bennett et al. 2003; Spergel et al. 2003).

To evaluate the molecular weight μ_e appearing in eqns. (3) and (6) we assume that the accreting plasma is a zero-metallicity, primordial gas, for which

$$\mu_e = \frac{1}{1 - Y/2}. \quad (22)$$

We take the primordial helium abundance Y to be $Y \approx 0.25$ (Cyburt, Fields & Olive 2003).

For the concordance model, the age of the universe is $T = t(0) = 13.7$ Gyr, while the age at redshift $z = 6.43$, the highest known quasar redshift, (SDSS 1148+525; Fan et al. 2003), is only $t_{\text{QSO}} = 0.87$ Gyr. Hence t_{QSO} is the upper limit to the time available for accretion to occur onto the initial seed black hole that powers this quasar. In fact, assuming that the black hole seed forms from the collapse of a first generation, Pop III star at redshift $z \lesssim 40$ and $t \gtrsim t(40) = 0.067$ Gyr, the available time for accretion is reduced to $t_{\text{accrete}} \lesssim 0.80$ Gyr. We note that the stellar evolution (hydrogen burning) lifetime of a massive Pop III progenitor, $t_{\text{evo}} \sim 0.003$ Gyr (Wagoner 1969; Onukai & Palla 2003), is much smaller than $t(40)$, so that the delay between stellar formation and collapse is of little consequence for determining the total time available for accretion growth. Most important, the exponential accretion growth timescale is given by eqns. (5), (6) and (22) to be

$$\tau_{\text{grwth}} = \frac{\epsilon_M}{1 - \epsilon_M} \frac{1}{\epsilon_L} \tau = 0.0394 \frac{(\epsilon_M/0.1)}{1 - \epsilon_M} \frac{1}{\epsilon_L} \text{ Gyr}, \quad (23)$$

which is considerably smaller than t_{accrete} . It is the availability of many exponential growth timescales from the time of black hole seed formation to the birth of a quasar that makes it possible for the black hole to grow from stellar to supermassive size by gas accretion (aided by mergers) in the early universe.

3. BLACK HOLE GROWTH AND SPINUP

Here we integrate the coupled mass and spin evolution equations (5) and (8) to study black hole growth and spin-up by gas accretion. In Fig. 1 we show the increase in mass by accretion at the Eddington limit ($\epsilon_L = 1$) as a function of time. In (a) the accretion disk is a standard thin disk with a “no torque boundary condition” at the ISCO (eqn. 14); in (b) the disk is an MHD disk as calculated by McKinney & Gammie (2004) (eqn. 15). We consider two different initial values for the black hole spin parameter, a/M . The solid lines show the case in which the initial black hole is nonrotating with $a/M = 0$. The dotted lines show the case in which the black hole to be spinning with $a/M = 0.75$. The latter is the value calculated for a black hole formed from the catastrophic collapse of a massive, radiation-dominated star spinning uniformly at the mass-shedding limit that has evolved to the onset of relativistic radial instability just prior to collapse (Shibata & Shapiro 2002; Shapiro & Shibata 2002; Shapiro 2004). For each disk model, the solid dots show the black hole growth that would occur if the accretion were maintained at the asymptotic, equilibrium spin and radiation efficiency from the beginning. Time is plotted in units of τ given by eqns. (6) and (22) for a zero-metallicity, cosmological abundance of H and He. The total duration plotted corresponds to the age difference between redshift $z_i = 40$ (the earliest plausible birthdate of a black hole from the collapse of a Pop III star) and redshift $z_f = 6.43$ (the highest known quasar redshift, corresponding to SDSS 1148+5251; Fan et al. 2003) in the adopted Λ CDM cosmology.

The key difference between the two disk models is that the standard thin disk drives the black hole to maximal spin and efficiency ($a/M = 1$ and $\epsilon_M = 0.42$; Bardeen 1970) while the MHD disk drives the black hole to spin equilibrium at lower values ($a/M \approx 0.95$ and $\epsilon_M \approx 0.19$). *A moderately lower radiation efficiency ϵ_M for Eddington-limited accretion results in substantially larger black hole growth.*

As is evident from the figure, apart from an initial transient lasting $\sim 0.1\tau \sim \tau_{\text{grwth}}$ (see figure inset), *the asymptotic evolution and final black hole mass amplification factor is little affected by the initial black hole spin.*

In Fig. 2 we show the spin evolution during Eddington-limited accretion for the cases shown in Fig. 1. The large figure plots the spin parameter versus time, while the inset plots the spin parameter as a function of black hole mass amplification. After the initial transient lasting $\sim 0.1\tau$, during which time the black hole grows by a factor of ~ 2 , the black hole spin and efficiency approach their asymptotic values. An analytic integration of eqn. (8) with eqn. (15), crudely holding ϵ_M fixed, yields an exponential spin-up timescale $\tau_{\text{spin}} \approx (1 - \epsilon_M)\tau_{\text{grwth}}/3.30$, which explains the rapid spin-up rate. This result also explains why the asymptotic black hole evolution and mass amplification are little affected by the initial black hole spin and depend only on the equilibrium spin rate




FIG. 1.— Black hole growth by accretion at the Eddington limit. In (a) the accretion disk is a standard thin disk with a “no torque boundary condition” at the ISCO, ignoring photon recapture; in (b) the disk is an MHD disk as calculated by McKinney & Gammie (2004). The solid lines show evolution starting from $a/M = 0$, while the dotted lines show evolution starting from $a/M = 0.75$. For each disk model, the solid dots show the mass amplification that would occur if the accretion were maintained at the asymptotic, equilibrium spin and radiation efficiency from the beginning. Time is plotted in units of $\tau = Mc^2/L_E = 0.45/\mu_e$ Gyr. The total duration shown corresponds to the age difference between redshift $z_i = 40$ (the earliest plausible birthdate of a black hole from the collapse of a Pop III star) and redshift $z_f = 6.43$ (the highest known quasar redshift, corresponding to 1148+5251) in a concordance Λ CDM cosmology.

of the accreting Kerr black hole. Thus, for all but the initial transient, it is adequate to adopt the equilibrium spin rate and use eqn. (16) with the corresponding constant efficiency $\epsilon_M[(a/M)_{\text{eq}}]$ to calculate the cosmological black hole mass amplification in lieu of a detailed integration of coupled evolution equations.

4. COSMOLOGICAL IMPLICATIONS

Next we consider some of the cosmological implications of the results of the previous section for the growth of SMBHs in the early universe.

In Fig. 3 we employ eqns. (16) and (21) to study the black hole mass amplification factor M_f/M_i as a function of the redshift z_i of the initial seed black hole. Here we plot the final amplification factor achieved by redshift $z_f = 6.43$, the highest known quasar redshift. Each solid curve is labelled by the adopted constant radiation efficiency, ϵ_M ; the luminosity is assumed to be the Eddington value ($\epsilon_L = 1$). The horizontal dashed lines indicate the range of amplification factors required for *accretion alone* to grow a supermassive black hole of mass $10^9 M_\odot$ from an initial seed black hole of mass $100 \leq M/M_\odot \leq 600$ formed from the collapse of a Pop III star. A mass of $10^9 M_\odot$ is the value inferred for typical quasars, including 1148+5251 (Fan et al. 2003). The lower the mass of the initial seed, the larger is the required mass amplification. The horizontal dotted lines indicate the required range of accretion-driven mass amplification assuming that *mergers* assist the growth process by accounting for an amplification of $f \sim 10^4$ in




FIG. 2.— Spin evolution during Eddington-limited accretion for the cases shown in Fig 1. Curves are labelled as in that figure. The large figure plots the spin parameter versus time, while the inset plots the spin parameter versus the black hole mass amplification factor.




FIG. 3.— Black hole accretion mass amplification M_f/M_i versus redshift z_i of the initial seed. Here we plot the amplification achieved by redshift $z_f = 6.43$, the highest known quasar redshift, corresponding to 1148+5251. Each solid curve is labelled by the adopted constant radiation efficiency, ϵ_M ; the luminosity is assumed to be the Eddington value ($\epsilon_L = 1$). The horizontal dashed lines bracket the range of amplification required for accretion alone to grow a seed black hole of mass $100 \leq M/M_\odot \leq 600$ formed from the collapse of a Pop III star to $10^9 M_\odot$. The horizontal dotted lines bracket the required accretion amplification range assuming that mergers account for a growth of 10^4 in black hole mass, the remainder being by gas accretion.

black hole mass, with the remaining amplification provided by gas accretion (Yoo & Miralda-Escudé 2004).

Some important cosmological implications can be inferred from Fig 3. Consider a seed black hole that forms sometime after redshift $z_i \lesssim 40$, by which time the earliest stars have formed and collapsed. In the absence of mergers, steady accretion cannot by itself achieve the

required growth to explain quasars at $z_f = 6.43$ unless the efficiency satisfies $\epsilon_M \lesssim 0.13$. For disk accretion, this constraint requires the steady-state black hole spin parameter to be below $a/M \approx 0.83$. This value is below (1) the maximal black hole spin $a/M = 1$, the asymptotic equilibrium value for a standard thin disk; (2) $a/M = 0.998$ (for which $\epsilon_M \approx 0.32$), the equilibrium value of a standard thin disk accounting for the recapture of some of the emitted photons by the black hole (Thorne 1974); and (3) $a/M \approx 0.95$, the equilibrium value of a typical MHD disk. Therefore, it is likely that *mergers are required to assist accretion to achieve black hole growth to supermassive size by $z_f = 6.43$* . As may be inferred from the figure, this conclusion holds even in the (unlikely) event that the black hole seed forms much earlier than $z_i \approx 40$.

Monte Carlo simulations by Yoo & Miralda-Escudé (2004) of hierarchical CDM halo mergers, accompanied by mergers of their central black holes, suggest that black hole mass amplification factors of $\sim 10^4$ are achieved via mergers by $z_f = 6.43$. This result implies that $f = f_1 f_2 \dots f_N \sim 10^4$ in eqn. (17). While other simulations predict lower growth factors due to mergers, we adopt 10^4 to be conservative, noting that our conclusions below are strengthened if the smaller factors turn out to be correct. As eqn. (17) shows, eqn. (16) can be used to estimate the cumulative mass amplification by gas accretion, even when steady growth by accretion is interrupted and enhanced by stochastic black hole mergers. Minor mergers of massive black holes with smaller holes tend to spin-down the massive hole (Hughes & Blandford 2003; Gammie, Shapiro & McKinney 2004). Major mergers of two black holes of comparable mass following binary inspiral drive the merged remnant to $a/m \gtrsim 0.7$. [See, e.g., Baumgarte & Shapiro 2003 for a review of binary black hole coalescence, and references; see Gammie, Shapiro & McKinney 2004 for a general discussion of black hole spin-up and spin-down mechanisms, and references]. In either case, after a short transient epoch, accretion will drive the merged remnant to the disk accretion equilibrium spin rate and corresponding mass efficiency (see Section 2), as assumed in employing eqn. (16).

Fig 3. indicates that for black hole seeds that arise from collapsed Pop III stars, merger-assisted mass amplification to SMBH status is easily achieved for typical MHD accretion disks, marginally possible for a standard thin disk in spin equilibrium accounting for photon recapture, but not possible for a standard thin disk that drives the black hole to maximal rotation. However, *if the black hole seed is less than $600 M_\odot$, the standard thin disk appears to be ruled out*: the required amplification places the lower dotted line in the plot above the curve for $\epsilon_M = 0.32$, corresponding to $a/M = 0.998$, the value for a standard thin disk in spin equilibrium, accounting for photon recapture. These conclusions again also hold even if the black seed forms much earlier than $z_i \approx 40$, while they may be tightened if the seed forms later. In fact, it may be likely that the seed forms later, at $z_i \lesssim 40$, given that even $4 - \sigma$ peaks in the density perturbation spectrum for the progenitor halo of SDSS 1148+5251 do not collapse until $z \sim 30$ in the Λ CDM concordance cosmology (see, e.g. Figure 5 in Barkana & Loeb 2001).

Moreover, the potential wells of the earliest halos are quite shallow (~ 1 km/s) and may not be able to retain enough gas to form stars. Nevertheless, the effect of altering the date of birth of the BH seed is not very great unless $z_i \lesssim 20 - 25$, as is evident from Fig. 3.

It is significant that *the range of equilibrium accretion disk radiation efficiencies required to achieve the necessary growth of a black hole seed to supermassive size by $z_f = 6.43$ is consistent with the values inferred observationally for R , the ratio of the QSO plus AGN luminosity density to the mass density of SMBHs in nearby galaxies: $\epsilon_M \geq R \sim 0.1 - 0.2$ (Soltan 1982; Yu & Tremaine 2002; Elvis et al. 2002).* This consistency supports the notion that accretion plays an important role in SMBH growth and is responsible for the acquisition of the bulk of the final mass of a SMBH. *The range of ϵ_M inferred from R favors accretion disk models that drive the black hole to spin equilibrium in the range $0.7 \lesssim a/M \lesssim 0.95$, well below maximal rotation and consistent with the values found in recent simulations of relativistic MHD accretion disks.*

How are our conclusions altered if (when) a quasar is discovered at a higher redshift, $z_{\text{QSO}} > 6.43$? We anticipate this possibility in Fig. 4, where we solve eqn. (16) to plot the mass efficiency ϵ_M required to build a supermassive black hole from a seed as a function of the host quasar redshift, z_{QSO} . Here we again assume that a seed black hole of mass $100 \leq M/M_\odot \leq 600$, formed from the collapse of a Pop III star at $z_i \gg z_{\text{QSO}}$, grows by z_{QSO} to $10^9 M_\odot$. While accreting, the black hole again is assumed to radiate at the Eddington luminosity, with $\epsilon_L = 1$. We set $t_i = 0$, $z_i = \infty$, so that the plotted radiation efficiency represents an *upper limit* to ϵ_M ; increasing t_i and lowering z_i decreases the required efficiency (but the decrease is small if, for example, we set $z_i \gtrsim 40$). The dashed lines indicate the range of efficiencies required for gas accretion alone to achieve the necessary growth. The dotted lines indicate the range required for gas accretion assuming that mergers assist the growth and account for an amplification of $f = 10^4$ in black hole mass. The horizontal solid lines bracket the range of efficiencies inferred observationally for R .

In the absence of mergers, the upper limit to the radiation efficiency required to build a supermassive black hole by $z_{\text{QSO}} \approx 6.43$ is $\epsilon_M \approx 0.14$, consistent with observational constraints implied by R , but inconsistent with the much higher values expected for a standard thin disk driven to spin equilibrium, and marginally inconsistent for an MHD accretion disk. The inconsistencies become worse as $z_{\text{QSO}} > 6.43$. In fact, the upper limit to ϵ_M falls below 0.1 for $z_{\text{QSO}} \gtrsim 8$, which would force ϵ_M to lie below the observationally inferred range for R and would therefore be difficult to understand.

When mergers are included, the upper limit to the efficiency required to build a SMBH by $z_{\text{QSO}} \approx 6.43$ increases to $\epsilon_M \sim 0.30$ (i.e., any black hole seed born at a finite redshift, $\infty > z_i > z_{\text{QSO}}$, must accrete at a lower efficiency than 0.30 to reach $10^9 M_\odot$). This upper limit is roughly consistent with the observational constraint on $\epsilon_M \geq R$ and with theoretical values for accretion from an MHD disk in spin equilibrium, but only marginally consistent for accretion from a standard thin disk in spin equilibrium accounting for photon recapture, and incon-




FIG. 4.— The radiation efficiency ϵ_M required to build a SMBH versus the redshift z_{QSO} of the host quasar. The seed black hole forms with a mass $100 \leq M/M_\odot \leq 600$ from the collapse of a Pop III star at $z_i \gg z_{\text{QSO}}$ and grows by z_{QSO} to $10^9 M_\odot$. Accretion occurs at the Eddington luminosity. The dashed lines bracket the range of efficiencies required for accretion alone to achieve the necessary growth. The dotted lines bracket the range required for accretion assuming that mergers account for a growth of 10^4 in black hole mass. The horizontal solid lines bracket the range of efficiencies inferred from R , the observed ratio of the QSO plus AGN luminosity density to the mass density of local SMBHs.

sistent for accretion from a standard thin disk that drives the black hole to maximal rotation. *Should a quasar be discovered at $z_{\text{QSO}} > 6.43$, it would appear that accretion from a standard thin disk will be ruled out:* the upper limit will fall below $\epsilon_M = 0.32$, the value for a standard thin disk in spin equilibrium accounting for photon recapture. *Should a quasar be discovered at $z_{\text{QSO}} \gtrsim 10$, ϵ_M would fall below 0.19 and the results would be difficult to reconcile with accretion from a typical MHD disk as modeled in recent simulations.* These critical values of z_{QSO} are all smaller if the initial black hole seed is smaller than $600 M_\odot$, as the top curve, which sets the limit, is lowered in the figure. Finally, should a quasar be discovered (perhaps unexpectedly) at $z_{\text{QSO}} \gtrsim 18$, the upper limit to ϵ_M would drop below the observationally inferred value 0.1 for all $z_i \geq z_{\text{QSO}}$. This drop would not be easily explained by any of these models of accretion, and might require super-Eddington accretion with $\epsilon_L > 1$ to keep $\epsilon_M \gtrsim 0.1$. Alternatively, the value of R , observed for local host galaxies, might be significantly smaller at high redshift, which would relax the constraint on ϵ_M . But achieving the inferred lower radiation efficiencies might then require ADAF disks or spherical accretion.

5. SUMMARY

We have explored the evolution of black hole mass and spin by gas accretion in the early universe. We have illustrated how for Eddington-limited accretion, the growth of a SMBH depends sensitively on the radiative efficiency, ϵ_M . For disk accretion, the mean efficiency is determined by the equilibrium black hole spin, which in turn depends on the torques acting on the gas near the black hole horizon. We have explored the consequences of the as-

sumptions that seed black holes are the remnants of collapsed Pop III stars that may form as early as $z \lesssim 40$ and can grow to $\sim 10^9 M_\odot$ by $z_{\text{QSO}} = 6.43$, the highest redshift discovered to date, corresponding to QSO SDSS 1148+5251. Allowing for growth both by accretion and mergers, simple theory suggests that the required mass amplification is possible provided the radiation efficiency satisfies $\epsilon_M \lesssim 0.2$, with the upper limit decreasing should a quasar be discovered at higher redshift $z_{\text{QSO}} > 6.43$. The inferred efficiency is consistent with the observed ratio R of the QSO plus AGN radiation density to the mass density of SMBHs in local galaxies, which suggests that an appreciable fraction of the mass of the final black hole is acquired by disk accretion, rather than by mergers. The inferred efficiency favors MHD accretion disk models that exert non-zero torques on the gas at the inner edge of the disk. These disks ultimately drive the black hole to spin equilibrium at $a/M \sim 0.95$, substantially below the maximum spin allowed for a Kerr black hole.

We have made a number of simplifying assumptions in an effort to provide a concrete calculation that illustrates how the SMBH initial seed, its cosmological growth, and accretion disk models may all be constrained by the existence of quasars at high redshift. Our conclusions are tentative, as there exist many caveats. For example, if the progenitors of black hole seeds are SMSs with $M \gg 10^3 M_\odot$ instead of Pop III stars, many of the constraints on the radiation efficiency ϵ_M , and the associated range of expected black hole spins, may have to be relaxed. On the other hand, if the progenitors have masses $\lesssim 100 M_\odot$ the constraints favoring low efficiencies $\epsilon_M \lesssim 0.2$ characterizing MHD accretion disks in spin equilibrium are strengthened. If the accretion luminosity is super-Eddington with $\epsilon_L > 1$, then the upper limit for ϵ_M increases (see eqn. 16), relaxing the constraints which favor an MHD disk over a standard thin disk. Super-Eddington accretion is possible theoretically (Ruszkowski & Begelman 2003). However, the Sloan Digital Sky Survey of 12,698 broad-line quasars in the redshift interval $0.1 \leq z \leq 2.1$ supports the Eddington value as a physical limit (McLure & Dunlop 2004). Moreover, given the observed value for R , a higher limit for ϵ_M would imply that a large fraction of the emitted radiation lies outside the optical bandwidths included in the determination of R , which seems unlikely. Finally, mergers may lead to black hole mass amplification factors smaller than $f \sim 10^4$, the typical value found in the Monte-Carlo calculations of Yoo & Miralde-Escudé, and the value assumed here. A smaller merger amplification factor requires larger accretion-driven amplification in order for a black hole seed to reach SMBH size by $z_{\text{QSO}} = 6.43$ (see eqn. 17). This would strengthen the constraints set by SMBH accretion growth on the upper limit to ϵ_M , although the efficiency must still be consis-

tent with R if the bolometric accretion luminosity resides mostly in the optical wavebands included in R and the value of R measured for local galaxies also applies at higher redshift. Of course, the peak contribution to the value of R arises from the average behavior of accreting BHs with $z \sim 3$; this ratio does not necessarily apply to individual objects like SDSS 1148+5251, nor need it apply to SMBHs at $z \approx 6$.

We have assumed that the mass of SDSS 1148+5251 is $\sim 10^9 M_\odot$; a lower value would relax many of our constraints, which a higher value would strengthen them. If the flux from this source were amplified by gravitational lensing, or beaming, then a lower mass estimate would be appropriate. However, no multiple images have been seen (Richards et al. 2004) and it has been shown that high amplification without at least two images is very improbable (Keeton et al. 2004). Strong beaming also seems unlikely since it would reduce the line/continuum ratio (Haiman & Cen 2002), which is not observed (Willott et al. 2003). In fact, assuming that the quasar emits at the Eddington luminosity gives a mass of $4.6 \times 10^9 M_\odot$ (Fan et al. 2003; Haiman 2004), and this higher value strengthens our conclusions somewhat.

Isolating the accretion growth of a seed black hole in the early universe from the hole's full dynamical history and environment does not allow us to account for other important correlations that provide clues to the formation of SMBHs. Such correlations include the SMBH mass vs. bulge luminosity relation, $M_{\text{BH}} \propto L_{\text{bulge}}$ (Kormendy & Richstone 1995), and the SMBH mass vs. velocity dispersion relation, $M_{\text{BH}} \propto v_c^4$ (Gebhardt et al. 2000; Ferrarese & Merritt 2000; Tremaine et al. 2002), inferred for nearby host galaxies. Only by performing detailed simulations that track the formation and growth of SMBHs in a cosmological setting governed by hierarchical halo mergers, black hole mergers, gas settling, star formation, and feedback can these correlations be reliably reproduced. We look forward to the next generation of simulations that incorporate the recent results of relativistic MHD accretion onto black holes, since, as demonstrated here, the outcome of these global simulations may depend sensitively on the local physics of such accretion flows.

It is a pleasure to thank T. Abel, C. Gammie, J. McKinney, J. Miraldi-Escudé, and Q. Yu for valuable discussions. We also thank the referee for a critical reading of the manuscript and valuable comments. A portion of this work was performed during the summer of 2004 at the Aspen Center for Physics, whose hospitality is gratefully acknowledged. This work was supported in part by NSF Grants PHY-0205155 and PHY-0345151 and NASA Grant NNG04GK54G at the University of Illinois at Urbana-Champaign.

REFERENCES

- Abel, T., Bryan, G. L., & Norman, M. L. 2000, *ApJ*, 540, 39
 ———. 2002, *Science*, 295, 93
 Balberg, S., & Shapiro, S. L. 2002, *Phys. Rev. Lett.*, 88, 101, 301
 Balbus, S. A., & Hawley, J. F. 1991, *ApJ*, 376, 214
 Bardeen, J. M. 1970, *Nature*, 226, 64
 Bardeen, J. M., & Petterson, J. A. 1975, *ApJ*, 195, L65
 Barkana, R., & Loeb, A. 2001, *Phys. Reports*, 349/2, 125
 Baumgarte, T. W., & Shapiro, S. L. 1999, *ApJ*, 527, L5
 ———. 2003, *Phys. Reports*, 376/2, 41
 Bennett, C. L. et al. 2003, *ApJS*, 148, 1
 Bromley, J. M., Somerville, R. S., & Fabian, A. C. 2004, *MNRAS*, 350, 456
 Bromm, V., & Loeb, A. 2003, *ApJ*, 596, 34
 Bromm, V., Coppi, P. S., & Larson, R. B. 1999, *ApJ*, 527, L5
 ———. 2002, *ApJ*, 564, 23
 Chandrasekhar, S. 1964a, *Phys. Rev. Lett.*, 12, 114 & 437

- . 1964b, *ApJ*, 140, 417
- Cyburrt, R. H., Fields, B. D., & Olive, K. A. 2003, *Phys. Lett. B* 567, 227
- De Villiers, J. P., & Hawley, J. F. 2003, *ApJ*, 589, 458
- De Villiers, J. P., Hawley, J. F., & Krolick, J. H. 2003, *ApJ*, 599, 1238
- De Villiers, J. P., Hawley, J. F., Krolick, J. H., & Hirose, S. 2004, *ApJ* submitted (astro-ph/0407092)
- Elvis, M., Risaliti, G., & Zamorani, G. 2002, *ApJ*, 565, L75
- Fan, X. et al. 2003, *AJ*, 125, 1649
- Favata, M., Hughes, S. A., & Holtz, D. E. 2004, *ApJ*, 607, L5
- Ferrarese, L., & Merritt, D. 2000, *ApJ*, 539, L9
- Figer, D. F. et al. 1998, *ApJ*, 506, 384
- Fishbone, L. G., & Moncrief, V. 1976, *ApJ*, 207, 962
- Fowler, W. A. 1964, *Rev. Mod. Phys.*, 36, 545, 1104E
- Gammie, C. F., Shapiro, S. L., & McKinney, J. C. 2004, *ApJ*, 602, 312
- Gebhardt, K., et al. 2000, *ApJ*, 539, L13
- Genzel, R., Eckart, A., Ott, T., & Eisenhauer, F. 1997, *MNRAS*, 291, 219
- Ghez, A. M., Becklin, E., Duchene, G., Hornstein, S., Morris, M., Salim, S., & Tanner, A. 2003, *Astron. Nachr./AN* 324, No. S1, 3.
- Ghez, A. M., Morris, M., Becklin, E. E., Tanner, A., & Kremenek, T. 2000, *Nature*, 407, 349
- Gnedin, O. Y. 2001, *Class. & Quant. Grav.*, 18, 3983
- Haiman, Z. 2004, *ApJ*, 613, 36
- Haiman, Z., & Cen, R. 2002, *ApJ*, 578, 702
- Haiman, Z., & Loeb, A. 2001, *ApJ*, 552, 459
- Haiman, Z., & Quataert, E. 2004, in *Supermassive Black Holes in the Distant Universe*, ed. A. J. Barger, (Dordrecht: Kluwer), in press (astro-ph/0403225).
- Haehnelt, M. G. 2004, in *Carnegie Observatories Astrophysics Series, Vol 1: Coevolution of Black Holes and Galaxies*, ed. L. C. Ho (Cambridge: Cambridge Univ. Press), p. 406
- Haehnelt, M. G., & Kauffmann, G. 2000, *MNRAS*, 318, L35
- Heger, A., Fryer, C. L., Woosley, S. E., Langer, N., & Hartmann, D. H. 2003, *ApJ*, 591, 288
- Ho, L. C. 1999, in *Observational Evidence for Black Holes in the Universe*, ed. S. K. Chakrabarti (Dordrecht: Kluwer), 157
- Hughes, S. A., & Blandford, R. D. 2003, *ApJ*, 585, L101
- Hut, P., & Rees, M. J. 1992, *MNRAS*, 259, 27p
- Keeton, C. R., Kuhlen, M., & Haiman, Z. 2004, *ApJ* submitted, (astro-ph/0405143)
- Kormendy, J., & Richstone, R. D. 1995, *Ann. Rev. Astr. & Ap.*, 33, 581
- Macchetto, F. D. 1999, in *Towards a New Millennium in Galaxy Morphology*, ed. D. L. Block, I. Puerari, A. Stockton & D. Ferreira (Dordrecht: Kluwer); *Astrophys.Space Sci* 269, 269
- Madau, P., & Rees, M. 2001, *ApJ*, 551, L27
- Madau, P., & Quataert, E. 2004, *ApJ*, 606, L17
- McKinney, J. C., & Gammie, C. F. 2004, *ApJ*, 611, 977
- McLure, R. J., & Dunlop, J. S. 2004, in *Proceedings of Multi-wavelength AGN Surveys*, Cozumel, 2003, in press (astro-ph/0405393)
- Meritt, D., Milosavljevic, M., Favata, M., Hughes, S. A., & Holz, D. E. 2004, *ApJ*, 607, L9
- Natarajan, P., & Pringle, J. E. 1998, *ApJ*, 506, L97
- Novikov, I. D., & Thorne, K. S. 1973, in *Black Holes*, Les Houches 1972, ed. C. DeWitt & B. DeWitt (Gordon & Breach: New York), p. 343
- Onukai, K., & Palla, F. 2003, *ApJ*, 589
- Ostriker, J. P. 2000, *Phys. Rev. Lett.*, 84, 5258
- Pringle, J. E., & Rees, M. J. 1972, *A & A*, 29, 179
- Norman, M. L. 2004, talk presented at Aspen Summer Workshop on The Formation of Supermassive Black Holes, June 2004, Aspen Center for Physics
- Rees, M. J. 1978, *Nature*, 275, 516
- . 1984, *Ann.Rev.Astr.&Ap* 22, 471
- . 1998, in *Black Holes and Relativistic Stars*, ed. R. M. Wald (Chicago: Chicago Univ. Press), 79
- . 2001, in *Black Holes in Binaries and Galactic Nuclei*, ed. L. Kaper, E. P. J. van den Heuvel, & P. A. Woudt (New York: Springer-Verlag), 351
- Richards, G. T., et al. 2004, *AJ*, 127, 1305
- Richstone, D., et al. 1998, *Science*, 395, A14
- Ruszkowski, M., & Begelman, M. C. 2003, *ApJ*, 586, 384
- Schödel, R., et al. 2002, *Nature*, 419, 694
- Shapiro, S. L. 2004, in *Carnegie Observatories Astrophysics Series, Vol 1: Coevolution of Black Holes and Galaxies*, ed. L. C. Ho (Cambridge: Cambridge Univ. Press), p. 103
- , 2004, *ApJ*, 610, 913
- Shapiro, S. L., & Shibata, M. 2002, *ApJ*, 577, 904
- Shapiro, S. L., & Teukolsky, S. A. 1979, *ApJ*, 234, L177
- , 1983, *Black Holes, White Dwarfs, and Neutron Stars: The Physics of Compact Objects* (New York: Wiley)
- Shibata, M., & Shapiro, S. L. 2002, *ApJ*, 527, L39
- Soltan, A. 1982, *MNRAS*, 200, 115
- Spergel, D. N. et al. 2003, *ApJS*, 148, 175
- Thorne, K. S. 1974, *ApJ*, 191, 507
- Tremaine, S., et al. 2002, *ApJ*, 574, 740
- Volonteri, M., Haardt, F., & Madau, P. 2003, *ApJ*, 582, 559
- Volonteri, M., & Madau, P., Quataert, E., & Rees, M. J. 2004, *ApJ*, in press (astro-ph/0410342)
- Wagoner, R. V. 1969, *Ann. Rev. Astron. & Ap.*, 7, 553
- Willott, C. J., McLure, R. J., & Jarvis, M. J. 2003, *ApJ*, 587, L15
- Yishida, N., Abel, T., Hernquist, L., & Sugiyama, N. 2003, 592, 645
- Yoo, J., & Miralde-Escudé, J. 2004, *ApJ Letters*, in press (astro-ph/0406217)
- Yu, Q., & Tremaine, S. 2002, *MNRAS*, 335, 965